



The Smallest Eigenvalue of a Large Dimensional Wishart Matrix

Author(s): Jack W. Silverstein

Source: *The Annals of Probability*, Vol. 13, No. 4, (Nov., 1985), pp. 1364-1368

Published by: Institute of Mathematical Statistics

Stable URL: <http://www.jstor.org/stable/2244186>

Accessed: 19/05/2008 08:03

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ims>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We enable the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.

THE SMALLEST EIGENVALUE OF A LARGE DIMENSIONAL WISHART MATRIX

BY JACK W. SILVERSTEIN¹

Weizmann Institute of Science

For positive integers s, n let $M_s = (1/s)V_s V_s^T$, where V_s is an $n \times s$ matrix composed of i.i.d. $N(0, 1)$ random variables. Assume $n = n(s)$ and $n/s \rightarrow y \in (0, 1)$ as $s \rightarrow \infty$. Then it is shown that the smallest eigenvalue of M_s converges almost surely to $(1 - \sqrt{y})^2$ as $s \rightarrow \infty$.

For each $s = 1, 2, \dots$ let $n = n(s)$ be a positive integer such that $n/s \rightarrow y > 0$ as $s \rightarrow \infty$. Let V_s be an $n \times s$ matrix whose entries are i.i.d. $N(0, 1)$ random variables and let $M_s = (1/s)V_s V_s^T$. The random matrix $V_s V_s^T$ is commonly referred to as the Wishart matrix $W(I_n, s)$.

It is well known [Marčenko and Pastur (1967), Wachter (1978)] that the empirical distribution function F_s of the eigenvalues of M_s [$F_s(x) \equiv (1/n) \times$ (number of eigenvalues of $M_s \leq x$)] converges almost surely as $s \rightarrow \infty$ to a nonrandom probability distribution function F_y having a density with positive support on $[(1 - \sqrt{y})^2, (1 + \sqrt{y})^2]$, and when $y > 1$, F_y yields additional mass on $\{0\}$. It is also known [Geman (1980)] that the maximum eigenvalue $\lambda_{\max}^{(s)}$ of M_s converges a.s. to $(1 + \sqrt{y})^2$ as $s \rightarrow \infty$. [The statement of this result in Geman (1980) has all the M_s constructed from one doubly infinite array of i.i.d. random variables. However, it is obvious from the proof that no relation on the entries of V_s for different s is needed.] These results are established under assumptions more general on the entries of V_s than Gaussian distributed, involving conditions on the moments of these random variables.

The present paper will prove the following

THEOREM. For $y < 1$ the smallest eigenvalue $\lambda_{\min}^{(s)}$ of M_s converges a.s. to $(1 - \sqrt{y})^2$ as $s \rightarrow \infty$.

The proof relies on Geršgorin's theorem [Geršgorin (1931)] which states: Each eigenvalue of an $n \times n$ complex matrix $A = (a_{ij})$ lies in at least one of the disks

$$|z - a_{jj}| \leq \sum_{i \neq j} |a_{ij}|, \quad j = 1, 2, \dots, n,$$

in the complex plane.

Geršgorin's theorem will be applied to a tridiagonal matrix orthogonally similar to M_s . This result is relevant to areas in multivariate statistics, for example regression or tests using the central multivariate F matrix, where the

Received May 1984; revised November 1984.

¹On leave from North Carolina State University. Supported under NSF Grant MCS-8101703A01. AMS 1980 subject classifications. Primary 60F15; secondary 62H99.

Key words and phrases. Smallest eigenvalue of random matrix, Geršgorin's theorem, χ^2 distribution.

boundedness of the largest eigenvalue of M_s^{-1} , namely $[\lambda_{\min}(s)]^{-1}$, is needed. The truth of the theorem for non-Wishart matrices would also be important. However, as will be seen, the proof relies strongly on the variables being normal, so a different method appears to be necessary for more general sample covariance matrices.

PROOF OF THE THEOREM. Since F_y has positive support to the right of $(1 - \sqrt{y})^2$ we immediately have

$$(1) \quad \limsup_{s \rightarrow \infty} \lambda_{\min}^{(s)} \leq (1 - \sqrt{y})^2 \quad \text{a.s.}$$

Assume s is sufficiently large so that $n < s$. Let O_s^1 be $s \times s$ orthogonal, its first column being the normalization of the first row of V_s , the remaining columns independent of the rest of V_s . The columns of O_s^1 can be constructed, for example, by performing the Gram-Schmidt orthonormalization process to the first row of V_s , together with $s - 1$ linearly independent nonrandom s -dimensional vectors. We have that $V_s^1 \equiv V_s O_s^1$ is such that its first row is $(X_s, 0, 0, \dots, 0)$, where X_s^2 is $\chi^2(s)$, $X_s \geq 0$, and the remaining rows are again made up of i.i.d. $N(0, 1)$ random variables. (It will also follow that X_s is independent of the remaining elements of V_s^1 but this fact will not be needed.)

Let O_n^1 be $n \times n$ orthogonal of the form

$$\begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & O_{n-1}^1 & \\ 0 & & & \end{pmatrix},$$

where O_{n-1}^1 is orthogonal, its first row being the normalization of $\{(V_s^1)_{j1}\}_{j=2}^n$ (as a vector in \mathbb{R}^{n-1}), the rest independent of V_s^1 . Then $V_s^2 \equiv O_n^1 V_s^1$ is of the form

$$\begin{pmatrix} X_s & 0 & \cdots & 0 \\ Y_{n-1} & & & \\ 0 & & & \\ \vdots & & W_{n-1, s-1} & \\ 0 & & & \end{pmatrix},$$

where Y_{n-1}^2 is $\chi^2(n - 1)$, $Y_{n-1} \geq 0$ and $W_{n-1, s-1}$ is $(n - 1) \times (s - 1)$, made up of i.i.d. $N(0, 1)$ random variables.

We then multiply V_s^2 on the right by an $s \times s$ orthogonal matrix O_s^2 of the form

$$\begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & & & & \\ 0 & & & & \\ \vdots & & O_{s-1}^2 & & \\ 0 & & & & \end{pmatrix}.$$

where the first column of O_{s-1}^2 is the normalization of the first row of $W_{n-1, s-1}$,

and then multiply $V_s^2 O_s^2$ on the left by an appropriate $n \times n$ orthogonal matrix, and so on. In the end we will have the existence of two orthogonal matrices O_n and O_s such that

$$O_n V_s O_s = \begin{pmatrix} X_s & 0 & 0 & 0 & & \cdots & & & & 0 \\ Y_{n-1} & X_{s-1} & 0 & 0 & & \cdots & & & & 0 \\ 0 & Y_{n-2} & X_{s-2} & 0 & & \cdots & & & & 0 \\ 0 & 0 & \vdots & \vdots & & \cdots & & & & 0 \\ \vdots & \vdots & & & & & & & & \vdots \\ 0 & 0 & 0 & 0 & \cdots & 0 & Y_1 & X_{s-(n-1)} & 0 & \cdots & 0 \end{pmatrix},$$

where X_i^2 is $\chi^2(i)$, $X_i \geq 0$, and Y_j^2 is $\chi^2(j)$, $Y_j \geq 0$. The fact that these random variables are independent will not be needed.

It follows that M_s is orthogonally similar to a tridiagonal matrix, the first and last rows being, respectively,

$$(1/s)(X_s^2, X_s Y_{n-1}, 0, \dots, 0),$$

$$(1/s)(0, 0, \dots, 0, X_{s-n+2} Y_1, Y_1^2 + X_{s-n+1}^2),$$

while the three nonzero elements in the $j + 1$ st row ($j = 1, 2, \dots, n - 2$) are

$$(1/s)(X_{s-j+1} Y_{n-j}, Y_{n-j}^2 + X_{s-j}^2, X_{s-j} Y_{n-j-1}).$$

By Geršgorin's theorem we have that

$$(2) \quad \lambda_{\min}^{(s)} \geq \min \left[(1/s)(X_s^2 - X_s Y_{n-1}), (1/s)(Y_1^2 + X_{s-n+1}^2 - X_{s-n+2} Y_1), \right. \\ \left. \min_{j \leq n-2} (1/s)(Y_{n-j}^2 + X_{s-j}^2 - (X_{s-j+1} Y_{n-j} + X_{s-j} Y_{n-j-1})) \right].$$

We have $\chi^2(1)/m \rightarrow_{a.s.} 0$ and $\chi^2(m)/m \rightarrow_{a.s.} 1$ as $m \rightarrow \infty$. Since $s/n \rightarrow y \in (0, 1)$ as $s \rightarrow \infty$ we have

$$(1/s)(X_s^2 - X_s Y_{n-1}) \rightarrow_{a.s.} 1 - \sqrt{y},$$

$$(1/s)(Y_1^2 + X_{s-n+1}^2 - X_{s-n+2} Y_1) \rightarrow_{a.s.} 1 - y \quad \text{as } s \rightarrow \infty.$$

Notice $1 - y > 1 - \sqrt{y} > (1 - \sqrt{y})^2$.

Applying Markov's inequality to $P(\exp(t\chi^2(m) - tm) > \exp(t\epsilon))$ and $P(\exp(-t\chi^2(m) + tm) > \exp(t\epsilon))$ for sufficiently small $t > 0$, it is straightforward to show for any $\epsilon > 0$ the existence of an $a \in (0, 1)$ depending only on ϵ such that

$$P(|(\chi^2(m)/s) - (m/s)| > \epsilon) \leq 2a^s$$

for all $s > 0$ and all positive integers $m \leq s$.

Therefore we can apply Boole's inequality on $2n - 2$ ($\leq \text{constant} \cdot s$) events to conclude that for any $\epsilon > 0$

$$P\left(\max_{s-(n-2) \leq m \leq s} |(X_m^2/s) - m/s| > \epsilon \text{ or } \max_{m \leq n-1} |(Y_m^2/s) - m/s| > \epsilon \right)$$

is summable. Therefore

$$\max \left[\max_{s-(n-2) \leq m \leq s} |(X_m^2/s) - m/s|, \max_{m \leq n-1} |(Y_m^2/s) - m/s| \right] \rightarrow_{a.s.} 0 \quad \text{as } s \rightarrow \infty.$$

We have

$$\begin{aligned} A_j^s &\equiv \left| (1/s)(Y_{n-j}^2 + X_{s-j}^2 - (X_{s-j+1}Y_{n-j} + X_{s-j}Y_{n-j-1})) \right. \\ &\quad \left. - \left((n-j)/s + (s-j)/s - \left(\sqrt{(s-j+1)/s} \sqrt{(n-j)/s} \right. \right. \right. \\ &\quad \left. \left. \left. + \sqrt{(s-j)/s} \sqrt{(n-j-1)/s} \right) \right) \right| \\ &\leq \left| (Y_{n-j}^2/s) - (n-j)/s \right| + \left| (X_{s-j}^2/s) - (s-j)/s \right| \\ &\quad + \left| (X_{s-j+1}/\sqrt{s})(Y_{n-j}/\sqrt{s}) - \sqrt{(s-j+1)/s} \sqrt{(n-j)/s} \right| \\ &\quad + \left| (X_{s-j}/\sqrt{s})(Y_{n-j-1}/\sqrt{s}) - \sqrt{(s-j)/s} \sqrt{(n-j-1)/s} \right|. \end{aligned}$$

Using the inequality $|\underline{a}\underline{b} - ab| \leq |\underline{a}^2 - a^2|^{1/2} |\underline{b}^2 - b^2|^{1/2} + |a| |\underline{b}^2 - b^2|^{1/2} + |b| |\underline{a}^2 - a^2|^{1/2}$ for $a, b, \underline{a}, \underline{b}$ nonnegative, together with the fact that the nonrandom fractions making up A_j^s are bounded by 1, we conclude that

$$\max_{j \leq n-2} A_j^s \rightarrow_{a.s.} 0 \quad \text{as } s \rightarrow \infty.$$

The expression

$$\begin{aligned} &(n-j)/s + (s-j)/s - \left(\sqrt{(s-j+1)/s} \sqrt{(n-j)/s} \right. \\ &\quad \left. + \sqrt{(s-j)/s} \sqrt{(n-j-1)/s} \right) \end{aligned}$$

achieves its smallest value when $j = 1$, for which we get

$$\begin{aligned} &(n-1)/s + (s-1)/s - \left(\sqrt{(n-1)/s} + \sqrt{(s-1)/s} \sqrt{(n-2)/s} \right) \\ &\rightarrow y + 1 - 2\sqrt{y} = (1 - \sqrt{y})^2 \quad \text{as } s \rightarrow \infty. \end{aligned}$$

Therefore, from (2) we have

$$\liminf_{s \rightarrow \infty} \lambda_{\min}^{(s)} \geq (1 - \sqrt{y})^2 \quad \text{a.s.}$$

which, together with (1) gives us

$$\lim_{s \rightarrow \infty} \lambda_{\min}^{(s)} = (1 - \sqrt{y})^2 \quad \text{a.s.} \quad \square$$

We note that the above proof can easily be modified to show $\lambda_{\max}^{(s)} \rightarrow (1 + \sqrt{y})^2$ for all $y > 0$.

REFERENCES

[1] GEMAN, S. (1980). A limit theorem for the norm of random matrices. *Ann. Probab.* **8** 252-261.
 [2] GERŠGORIN, S. A. (1931). Über die Abgrenzung der Eigenwerte einer Matrix. *Izv. Akad. Nauk SSSR Ser. Fiz.-Mat.* **6** 749-754.

- [3] MARČENKO, V. A. and PASTUR, L. A. (1967). Distributions of eigenvalues of some sets of random matrices. *Math. USSR-Sb.* **1** 507–536.
- [4] WACHTER, K. W. (1978). The strong limits of random matrix spectra for sample matrices of independent elements. *Ann. Probab.* **6** 1–18.

JACK W. SILVERSTEIN
DEPARTMENT OF MATHEMATICS
BOX 8205
NORTH CAROLINA STATE UNIVERSITY
RALEIGH, NORTH CAROLINA 27695-8205