


**Exercise 1** *Polarization observable and measurement principle*

Consider the “measurement apparatus” (in the below figure) constituted of “an analyzer and a detector”. The incoming (initial) state of the photon is linearly polarized:

$$|\theta\rangle = \cos \theta |x\rangle + \sin \theta |y\rangle .$$



When the photodetector clicks we record  $+1$  and when it does not click we record  $-1$ . Thus the “polarization observable” is represented by the  $2 \times 2$  matrix

$$P_\alpha = (+1) |\alpha\rangle \langle\alpha| + (-1) |\alpha_\perp\rangle \langle\alpha_\perp|$$

where  $|\alpha\rangle = \cos \alpha |x\rangle + \sin \alpha |y\rangle$  and  $|\alpha_\perp\rangle = -\sin \alpha |x\rangle + \cos \alpha |y\rangle$  are the two vectors of the measurement basis. Note that the two orthogonal projectors of the measurement basis are  $\Pi_\alpha = |\alpha\rangle \langle\alpha|$  and  $\Pi_{\alpha_\perp} = |\alpha_\perp\rangle \langle\alpha_\perp|$ .


- 1) Show that  $\Pi_\alpha^2 = \Pi_\alpha$ ,  $\Pi_{\alpha_\perp}^2 = \Pi_{\alpha_\perp}$  and  $\Pi_\alpha \Pi_{\alpha_\perp} = \Pi_{\alpha_\perp} \Pi_\alpha = 0$ .
- 2) Check the following formulas:

$$\begin{aligned} |\langle\theta|\alpha\rangle|^2 &= \langle\theta|\Pi_\alpha|\theta\rangle, \\ |\langle\theta|\alpha_\perp\rangle|^2 &= \langle\theta|\Pi_{\alpha_\perp}|\theta\rangle \end{aligned}$$

- 3) Let  $p = \pm 1$  the random variable corresponding to the event click / no-click of the detector. Express  $\text{Prob}(p = \pm 1)$  with simple trigonometric functions and check that the two probabilities sum to one.
- 4) Deduce from 3)  $\mathbb{E}(p)$  and  $\text{Var}(p)$  and check that you find the same expressions by directly computing  $\langle\theta|P_\alpha|\theta\rangle$  and  $\langle\theta|P_\alpha^2|\theta\rangle - \langle\theta|P_\alpha|\theta\rangle^2$  in Dirac notation.

**Exercise 2** *Interferometer with an atom on the upper path*

Consider the following set-up where an atom may absorb the photon on the upper arm of the interferometer.



The Hilbert space of the photon is here  $\mathbb{C}^3$  with basis states

$$|H\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, |V\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, |\text{abs}\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

the semi-transparent and reflecting mirrors are modeled by the unitary matrices

$$S = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad R = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

and the “absorption-reemission” process<sup>1</sup> is modeled by the unitary matrix

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$$

Note that in Dirac notation

$$A = |H\rangle \langle \text{abs}| + |V\rangle \langle V| + |\text{abs}\rangle \langle H|$$

This models three possible transitions:  $A|H\rangle = |\text{abs}\rangle$  (absorption);  $A|\text{abs}\rangle = |H\rangle$  (emission); and  $A|V\rangle = |V\rangle$  (nothing happens).

- 1) Write down all matrices in Dirac notation and then compute the unitary operator  $U = SARS$  representing the total evolution process of this interferometer.
- 2) Given that the initial state is  $|H\rangle$ , what is the state after the second semi-transparent mirror? What are the probabilities of the following three events: click in  $D_1$ ; or click in  $D_2$ ; or no clicks in  $D_1$  nor  $D_2$ ? Verify the probabilities sum to 1.

<sup>1</sup>On the picture  $E_0$  and  $E_1$  are two energy levels of the atom corresponding to ground state and excited state; but you can ignore this aspect in this problem.

- 3) Suppose the photon-atom interaction is not absorption-reemission but some other process modeled by a matrix. Which of the two following matrices would be legitimate in QM?

$$\begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \text{ or } \begin{pmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{2}} \end{pmatrix}$$

and why?

**Exercise 3** *Entanglement for two quantum bits*

Consider two quantum bits in the state:

$$|\Psi\rangle = \frac{1}{\sqrt{3}} (|0\rangle \otimes |0\rangle + |1\rangle \otimes |1\rangle + i |1\rangle \otimes |0\rangle)$$

where  $|0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $|1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  are the canonical orthonormal basis of  $\mathbb{C}^2$ .

- 1) Write this state in 4-component-form as a column vector (use the conventions of class for the tensor product).
- 2) Prove that this state is “entangled” (in french ”intriquer”) in the sense that *it is impossible to express it in “product form”*

$$(\alpha |0\rangle + \beta |1\rangle) \otimes (\gamma |0\rangle + \delta |1\rangle)$$

for any  $\alpha, \beta, \gamma, \delta \in \mathbb{C}$ .