

In this exercise we illustrate a toy model description of the “orbital” state of a photon and its manipulation by mirrors. We suppose that a photon can travel only along the “vertical” and “horizontal” directions and represent its general state by a quantum bit

$$\alpha |H\rangle + \beta |V\rangle = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$$

where  $|H\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $|V\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ . Recall that  $\alpha, \beta \in \mathbb{C}$  and  $\alpha^* \alpha + \beta^* \beta = 1$ .




Figure 1: Possible preparation directions and states

- 1) Write down the “Bra” in Dirac and usual vector notations associated to the “Ket”  $\alpha |H\rangle + \beta |V\rangle$ .
- 2) Compute the scalar product (or Bracket) for the two kets  $\alpha |H\rangle + \beta |V\rangle$  and  $\gamma |H\rangle + \delta |V\rangle$  in Dirac and vector notations. In particular, check  $\langle H|V\rangle = \langle V|H\rangle = 0$  and  $\langle H|H\rangle = \langle V|V\rangle = 1$ .
- 3) A mirror like Figure 2 makes the transitions  $|H\rangle \rightarrow i|V\rangle$  and  $|V\rangle \rightarrow i|H\rangle$ . In quantum





Figure 2: Mirror operations (perfect reflection)

physics the mirror operation is described by a matrix  $R = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$ . Verify  $R^\dagger R = RR^\dagger =$

$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  where  $R = R^{\top,*}$  (transpose and complex conjugate).


A photon in the state  $\alpha |H\rangle + \beta |V\rangle$  is incident on the mirror and is reflected. Compute the output state  $R(\alpha |H\rangle + \beta |V\rangle)$  in Dirac and vector notations. Make a picture of the photon and mirror.

- 4) A semi-transparent mirror is described by a matrix  $S = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix}$ . Verify  $S^\dagger S = SS^\dagger = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ . Compute  $S|H\rangle$ ,  $S|V\rangle$  and  $S(\alpha|H\rangle + \beta|V\rangle)$ . Make pictures analogous to Figure 2 to get an intuition of these operations.
- 5) Consider the following experiment where  $D$  is a photo-detection.



Here we will assume  $\alpha$  and  $\beta$  are real. Compute the probability of detecting a photon in  $D$ .

- 6) Now we consider the following set-up which constitutes the Mach-Zehnder interferometer.



The incoming photon has the state  $|H\rangle$ . Compute the final state just after the second semi-transparent mirror. Then compute the probability of detecting the photon in  $D_1$  and  $D_2$ . What would you expect to find if the photons were “classical balls”?